

As directed, IODEX AR/18 is effective in hard water (up to 400 ppm as CaCO₃) and in 5% organic serum load against the following viruses, bacteria and pathogenic fungi: Polio 1*, Vaccinia*, Herpes Simplex Type 1*, Influenza A/2 Japan 305*, Hepatitis B and Canine Parvovirus*; bactericidal against Salmonella choleraesuis, Staphylococcus aureus, Pseudomonas aeruginosa, and E. Coli; is fungicidal against Trichophyton mentagrophytes (the athlete's foot fungus).

On pre-cleaned food contact surfaces, IODEX AR/18 is effective against Salmonella choleraesuis, Listeria monocytogenes, Escherichia coli, 0157:H7, Hemorrhagic.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NOT FOR RESIDENTIAL USE

The color of an IODEX AR/18 solution is proportional to the titratable iodine concentration. Prepare a fresh solution daily, or when there is a noticeable change in its rich amber color, or more often if the solution becomes diluted or soiled.

ONE-STEP CLEANING AND DISINFECTING:

PRIOR TO USE in Veterinary Clinics, Animal Handling Facilities, Kennels, Pet Shops and Zoos: Remove all animals and feeds from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and all surfaces to be treated. Empty all troughs, racks and other feeding and watering appliances. Then proceed as directed below.

TO CLEAN AND DISINFECT: Before applying disinfecting solution, food products and packaging materials must be removed from the room or carefully protected. Remove heavy soil deposits before applying the disinfecting solution. Using a solution of 3 ounces IODEX AR/18 per 5 gallons of warm water (68°F, 20° C or warmer), immerse objects in disinfecting solution, or apply with mop, sponge or brush. Where appropriate, use mechanical sprayer to produce coarse spray. Allow a 10 minute contact time with disinfecting solution. For efficacy against Canine Parvovirus, allow a 120 minute contact time; surface must remain wet. All food contact surfaces must be rinsed with potable water before reuse. Floors, walls and all other non-food contact surfaces do not require rinsing before reuse.

NOTE: When using IODEX AR/18 against the following viruses, bacteria and pathogenic fungi: Polio 1*, Vaccinia*, Herpes Simplex Type 1*, Influenza A/2 Japan 305*, Salmonella choleraesuis, Staphylococcus aureus, Pseudomonas aeruginosa, and E. Coli; Trichophyton mentagrophytes in a solution of cold water (as low as 41°F, 5°C), increase use to 6 oz. per 5 gallons of water.

BEFORE REHOUSING ANIMALS: Ventilate buildings, vehicles and other closed spaces. Do not house animals or use equipment until treatment absorbs, sets or dries. Thoroughly rinse all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with potable water before reuse.

SHOE BATH: Routine use of shoe bath sanitizers helps reduce the spread of disease causing organisms between poultry houses, farrowing houses, hog barns and other livestock buildings. Use 3 ounces of IODEX AR/18 per 1 gallon of water. Place sanitizing solution in shallow plastic or stainless steel pans inside all entrances to poultry and livestock buildings. Scrape shoes outside doorway and stand in shoe bath for 30 seconds before entering building. Replace shoe bath daily or more often if solution becomes soiled.

REMOVAL OF MILKSTONE AND HARD WATER DEPOSITS: Yellow color may appear on equipment after the first use of IODEX AR/18. This may indicate the presence of milkstone or hard water deposits. Remove these deposits by applying a solution containing equal parts of IODEX AR/18 and lukewarm water. Wait 2-3 minutes, brush and then rinse with potable water. Daily use of IODEX AR/18 will help prevent the formation of milkstone and hard water deposits.

SANITIZING POULTRY DRINKING WATER: Add 1 ounce of IODEX AR/18 per 10 gallons of drinking water. Regular use of this product in drinking water reduces the buildup of slime and mineral deposits in watering equipment. Do not use this product together with other water treatments such as medications.

SANITIZING PREVIOUSLY CLEANED FOOD CONTACT SURFACES: Sanitize previously cleaned and rinsed hard, non-porous surfaces such as glass, metal, plastic and porcelain with a solution containing 1 ounce of IODEX AR/18 per 5 gallons of water. Immerse equipment with sanitizing solution and allow 1 minute of contact time. Drain solution from equipment; do not rinse. IODEX AR/18 used at 1 ounce per 5 gallons of water contains 25 ppm titratable iodine and does not require a final rinse with potable water in accordance with Federal Food Additive Regulation 178.1010.

FOR INDUSTRIAL, INSTITUTIONAL AND AGRICULTURAL USE

IODEX AR/18

VIRUCIDE*, BACTERICIDE, FUNGICIDE, DEODORIZER

CONCENTRATED BROAD SPECTRUM IODOPHOR FOR USE AS A ONE-STEP CLEANER-DISINFECTANT AND NO-RINSE SANITIZER IN: Veterinary Clinics, Poultry Drinking Water, Sanitizing Commercial Eggs, Egg Processing Plants, Poultry Farms, Meat & Poultry Plants, Food Processing Plants, Kennels, Pet Shops & Zoos, Hand Sanitizing in Food Plants, Animal Handling Facilities

Active Ingredient:

Iodine* 1.75%

Inert Ingredients: 98.25%

Total 100.00%

*From alpha-(p-nonylphenyl)-omega-hydroxypoly (oxyethylene)-iodine complex

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

KEEP OUT OF REACH OF CHILDREN DANGER

CORROSIVE: Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and aquatic organisms. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit, and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

See side panel for FIRST AID.

EPA Reg. No. 4959-15-46269

EPA Est. No. 46269-DE-01

NET CONTENTS: 5 GALLONS

EXP. DATE: MM/YYYY



1500 EASTLAWN AVENUE • WILMINGTON, DE 19802

TEL: 302-761-2600 FAX 302-761-2611

SALES OFFICE : 1-800-424-2436

SANITIZING FOOD GRADE EGG SHELLS: Coarse spray previously cleaned food-grade egg shells with a solution containing 1 ounce of IODEX AR/18 per 5 gallons of water. Solution temperature should be warmer than the eggs, but not to exceed 130° F (53° C). Wet eggs thoroughly for 1 minute and allow to drain. A final potable water rinse is not required. Eggs should be reasonably dry before casing or breaking. Do not reuse solution to sanitize eggs.

HAND SANITIZING: Thoroughly wash and rinse hands before sanitizing. Dip or rinse hands in a solution containing 1 ounce of IODEX AR/18 per 5 gallons of water. IODEX AR/18 may be injected directly into wash or rinse water at a rate of 1 ounce per 5 gallons of water. A final potable water rinse is not required.

DEODORIZING: IODEX AR/18 destroys many odors as it sanitizes and disinfects. For general deodorant applications (garbage pails, refuse containers, etc.), swab surfaces to be treated with a solution containing 1 ounce IODEX AR/18 per gallon of water. Allow treated surface to adequately drain.

USE DILUTION TABLE:

IODEX AR/18 to WATER	TITRATABLE IODINE
1 oz. to 10 Gallons	12.5 ppm
1 oz. to 5 Gallons	25 ppm
3 oz. to 5 Gallons	75 ppm
1 oz. to 1 Gallon	125 ppm
6 oz. to 5 Gallons	150 ppm

FIRST AID

If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes. Then continue rinsing. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor for treatment advice. Have the person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If Inhaled: Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency treatment information. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency treatment information.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsions may be needed.

STORAGE AND DISPOSAL

Keep container closed when not in use. Do not store below 25°F or above 100°F for extended periods.

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse, or equivalent, promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure for two more times.