

This product is a versatile Disinfectant and Virucide for Veterinarian, Veterinary Practice, Animal Care, and Animal Laboratory applications.

**Sani-Plex CPV** for use in:

• Veterinary clinics, animal life science laboratories, animal research centers, animal quarantine areas, animal breeding facilities, kennels, dog/cat animal kennels, breeding establishments, zoos, operating rooms, washing areas, waiting rooms, examination rooms, and other animal care facilities.

For use on hard non-porous surfaces in animal care facilities such as floors, walls, cabinets, chairs, tables, cages, feeding and watering equipment, furniture, operating tables, metal surfaces, stainless steel, glazed porcelain, painted or sealed woodwork, glazed ceramic tile and plastic.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### PREPARATION OF DISINFECTION/FUNGICIDAL/VIRUCIDAL USE

**SOLUTION:** To disinfect hard inanimate surfaces such as walls, floors, table tops, apply 3 ounces of this product per 1 gallon of water, with a cloth, mop, sponge or mechanical spray device. When applied with a mechanical spray device, surfaces must be sprayed until thoroughly wetted. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray. Treated surfaces must remain wet for 10 minutes. Then let surface air dry. Prepare a fresh solution at least daily or sooner if use solution becomes visibly dirty.

**NOTE:** With spray applications cover or remove all food products.

### DISINFECTION

**DISINFECTION IN NON-MEDICAL FACILITIES:** Add 3 ounces of this product to one gallon of water, or equivalent dilution. Treated surfaces must remain wet for 10 minutes. At this use level, this product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum against:

Corynebacterium ammoniagenes	Listeria monocytogenes
Bordetella bronchiseptica	Salmonella enterica
Enterobacteriaceae w/beta lactamase resistance	Salmonella schottmuelleri
Enterobacter aerogenes	Salmonella typhi
Enterococcus faecalis	Serratia marcescens
Enterococcus faecalis	Shigella dysenteriae
Vancomycin Resistant	Staphylococcus aureus
Escherichia coli	Staphylococcus aureus-MRSA
Klebsiella pneumoniae	Staphylococcus aureus-VISA
	Streptococcus salivarius

**\*VIRUCIDAL PERFORMANCE:** At 3 ounces per one gallon use level or equivalent dilution, this product was evaluated in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard, non-porous environmental surfaces: Human Coronavirus, Herpes Simplex Type 1, Herpes Simplex virus type 2, Influenza A/Asian virus, Influenza A2/Japan virus.

**\*ANIMAL PREMISE VIRUCIDAL PERFORMANCE:** At 3 ounces per gallon use level, this product was evaluated in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard nonporous environmental surfaces:

Canine Adenovirus	Feline Picomavirus
Canine Coronavirus	Feline Rhinotracheitis
Canine Distemper Virus	Feline leukemia
Canine Parainfluenza Virus	Infectious Canine Hepatitis Virus
Feline Calicivirus	Murine Parainfluenza
Feline Infectious Peritonitis	virus type 1
Feline Panleukopenia	Pseudorabies Virus
	Rabies Virus

At 9 ounces per gallon use level, this product is found to be effective against the following viruses on hard, non-porous environmental surfaces in the presence of 5% serum with a 10 minute contact time: Canine Parvovirus-CPV, Minute virus of Mice-Parvovirus, Porcine Parvovirus.

**\*FOR CANINE PARVOVIRUS ACTIVITY:** This product is effective against Canine Parvovirus at 9 ounces per gallon. Follow "Veterinary Clinics/Animal Life Science Laboratory/Animal Care Facilities/Animal Research Centers/Animal Quarantine Areas/Kennels/Animal Breeding Facilities/Breeding and Grooming Establishment Disinfection Directions".

# SANI-PLEX CPV

Disinfectant, Virucide\* - With Organic Soil  
Tolerance for Veterinary Clinics, Kennels, Animal  
Care Facilities, Animal Life Science Laboratories  
and Breeding Establishments.

### ACTIVE INGREDIENTS:

Alkyl (C<sub>14</sub> 60%, C<sub>16</sub> 30%, C<sub>12</sub> 5%, C<sub>18</sub> 5%)  
dimethyl benzyl ammonium chloride . . . . . 1.60%

Alkyl (C<sub>12</sub> 68%, C<sub>14</sub> 32%)  
dimethyl ethylbenzyl ammonium chloride . . . . . 1.60%

**INERT INGREDIENTS:** 96.80%  
**TOTAL:** 100.00%

## KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

See right panel for additional Precautionary Statements.

### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Reg. No. 10324-140-46269

EPA Est. No. 46269-DE-01

NET CONTENTS: 5-55 Gallons

EXP. DATE: \_\_\_\_\_

Lot #: \_\_\_\_\_



QUIP LABORATORIES, INC.

1500 EASTLAWN AVENUE • WILMINGTON, DE 19802  
TEL : 800-424-2436 FAX: 302-761-2611

VETERINARY CLINICS / ANIMAL LIFE SCIENCE LABORATORY / ANIMAL CARE FACILITIES / ANIMAL RESEARCH CENTERS / ANIMAL QUARANTINE AREAS / KENNELS / ANIMAL BREEDING FACILITIES / BREEDING AND GROOMING ESTABLISHMENT DISINFECTION

**DIRECTIONS:** For cleaning and disinfecting the following hard non-porous surfaces: equipment, utensils, instruments, cages, kennels, stables, stalls, and catteries. Remove all animals and feeds from areas to be treated, animal transportation vehicles, crates etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 3 ounces per gallon of water, 9 ounces for Canine Parvo Virus, Minute virus of Mice-parvovirus, and Porcine Parvovirus, or equivalent dilution, for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other equipment that dispenses food or water with soap or detergent, and rinse with potable water before reuse.

**FUNGICIDAL PERFORMANCE:** At 3 ounces per one gallon use level or equivalent dilution, this product was evaluated in the presence of 5% serum with a 10 minute contact time and found to be effective against the following fungi on hard, non-porous surfaces: Trichophyton mentagrophytes, Aspergillus niger

**MOLD AND MILDEW PERFORMANCE:** Add 3 ounces per gallon of water to control the growth of mold and mildew, Aspergillus niger, and their odors on hard, non-porous surfaces. Thoroughly wet all treated surfaces completely. Treated surfaces must remain wet for 10 minutes. Let air dry. Repeat application weekly or when growth or odor reappear.

### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

**Keep Out of Reach of Children.** Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed, inhaled, or absorbed through the skin. Do not get into eyes, on skin, or on clothing. Avoid breathing vapor or spray mist. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, using tobacco or using the toilet. Remove contaminated clothing and washclothing before reuse.

### PHYSICAL OR CHEMICAL HAZARDS

Do not mix with oxidizers, anionic soaps and detergents.

Do not use or store near heat or open flame.

### ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Open dumping is prohibited. Store only in original container. Do not reuse empty container. If a leaky container must be contained within another, mark the outer container to identify the contents. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING DISPOSAL:** Non-refillable container. Do not reuse this container to hold materials other than pesticides or diluted pesticides. Offer for recycling if available or puncture and dispose in a sanitary landfill, or by other procedures approved by state and local authorities. If rinsate cannot be used, follow pesticide disposal instructions. If not triple rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state and federal regulations.

**RESIDUE REMOVAL INSTRUCTIONS:** Triple rinse container, or equivalent, promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store for later use or disposal. Repeat this procedure two more times.