

MB-10 Tablets® are a unique formulation of sodium chlorite that rapidly produces the biocidal agent chlorine dioxide when mixed with water.

*A 100 ppm use-solution of chlorine dioxide has been shown to be effective against: Vaccinia virus, Human Influenza A virus (Hong Kong), Porcine Respiratory and Reproductive Syndrome virus (PRRSV), Infectious bursal disease virus (IBDV), Marek's disease virus (MDV), Avian Influenza A (H3N2) virus, Porcine circovirus type 2, Canine Parvovirus, Hantavirus, Minute Virus of Mouse (Parvovirus) (MVM-p), Minute Virus of Mouse (Parvovirus) (MVM-i), Mouse Hepatitis Virus (MHV-A59), Mouse Hepatitis Virus (MHV-JHM), Mouse Parvovirus type 1 (MPV-1), Murine Parainfluenza virus type 1 (Sendai), Sialodacryoadenitis Virus (Coronavirus) (SDAV), and Theiler's Mouse Encephalomyelitis Virus (TMEV) after 10 minutes of contact and Foot and Mouth Disease Virus after 30 minutes of contact.

A 200 ppm use-solution of chlorine dioxide is a broad spectrum disinfectant effective against Hepatitis B Virus, Hepatitis C Virus, Newcastle Disease virus, Norovirus, Human coronavirus and the following gram negative and gram positive bacteria: Escherichia coli O157:H7, Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella enterica, Bordetella bronchiseptica, Corynebacterium bovis and Helicobacter pylori and Clostridium difficile (vegetative) after 10 minutes of contact. The 200 ppm use-solution is also effective against Staphylococcus aureus (MRSA) and Enterococcus faecalis (VRE) after 5 minutes of contact, tuberculocidal against Mycobacterium bovis after 5 minutes of contact at 20°C and virucidal against HIV-1 (the virus that causes AIDS) after 1 minute of contact.

MB-10 Tablets® can be used as a disinfectant / virucide* / tuberculocide in Hospitals, Medical, Dental, Industrial, Institutional and Manufacturing Facilities, Laboratory Animal Facilities, Clinical and Research Laboratories, Veterinary Clinics and Hospitals, Animal Rearing and Confinement Facilities, Animal Research Facilities and Laboratories, and other institutional/industrial applications that involve the housing of animals.

MB-10 Tablets® can also be used as a sterilant in manufacturing facilities, clinical laboratories, and Biosafety Level 3 and 4 (BSL-3 and BSL-4) facilities.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PREPARATION OF USE-SOLUTION:

25 ppm Chlorine Dioxide (for sanitization of hard, nonporous food contact surfaces).

In a clean plastic pail, place one (1) 1.5 gram **MB-10 Tablet**® for every gallon of clean potable water. Or, in a plastic pigmented spray bottle, place one (1) 1.5 gram **MB-10 Tablet**® for every gallon (4 liters) of clean potable water. Prepare in a well ventilated area. Wait 10 minutes for the 1.5 gram tab to completely dissolve. Once dissolved, this will yield a working solution of **25 ppm** of free chlorine dioxide. Once mixed, this solution should be stored in a tightly covered container and used within 7 days.

50 ppm Chlorine Dioxide (for sanitization of hard, nonporous food contact surfaces against Listeria monocytogenes).

In a clean plastic pail, place two (2) 1.5 gram **MB-10 Tablets**® for every gallon of clean potable water. Or, in a plastic pigmented spray bottle, place one (1) 1.5 gram **MB-10 Tablets**® for every two (2) quarts (or two (2) liters) of clean, potable water. Prepare in a well ventilated area. Wait 10 minutes for the 1.5 gram tab to completely dissolve. Once dissolved, this will yield a working solution of 50 ppm of free chlorine dioxide. Once mixed, this solution should be stored in a tightly covered container and used within 7 days.

100 ppm Chlorine Dioxide (for control of Foot and Mouth Disease Virus, Human Influenza A virus (Hong Kong), Avian Influenza A (H3N2) virus and animal viruses*).

In a clean plastic pail or plastic pigmented 1 gallon bottle, place either one (1) 6.0 gram **MB-10 Tablets**® or four (4) 1.5 gram **MB-10 Tablets**® for every gallon (4 liters) of clean potable water. Or, in a plastic pigmented spray bottle, place one (1) 1.5 gram **MB-10 Tablets**® for every quart (or liter) of clean, potable water. Prepare in a well ventilated area. Once dissolved, this will yield a working solution of **100 ppm** of free chlorine dioxide. Wait 10 minutes for the 1.5 gram tab and 15 minutes for the 6.0 gram tab to completely dissolve. Once mixed, this solution should be stored in a tightly covered container and used within 7 days.

200 ppm Chlorine Dioxide (for disinfection of hard, nonporous surfaces and instruments, and to kill tuberculosis bacteria, Human coronavirus, HIV, Hepatitis B virus, Hepatitis C virus, Newcastle Disease virus and Norovirus).

In a clean plastic pail or plastic pigmented 1 gallon bottle, place two (2) 6.0 gram **MB-10 Tablets**® or eight (8) 1.5 gram **MB-10 Tablets**® for every gallon (4 liters) of clean potable water. Or, in a plastic pigmented spray bottle, place one (1) 1.5 gram **MB-10 Tablets**® for every pint (or 500 ml) of clean, potable water. Prepare in a well ventilated area. Wait 10 minutes for the 1.5 gram tab and 15 minutes for the 6.0 gram tab to completely dissolve. Once dissolved, this will yield a working solution containing **200 ppm** of free chlorine dioxide. Once mixed, this solution should be stored in a tightly covered container and used within 7 days.

1000 ppm Chlorine Dioxide (for surface sterilization).

In a clean plastic pail or plastic pigmented 1 gallon bottle, place eight (8) 6.0 gram **MB-10 Tablets**® for every gallon of clean potable water or eight (8) 1.5 gram **MB-10 Tablets**® for every liter of clean potable water. Or, in a plastic pigmented spray bottle, place one (1) 6.0 gram **MB-10 Tablets**® for every pint (or 500 ml) of clean, potable water. Prepare in a well ventilated area. Wait 15 minutes for the 6.0 gram tab to completely dissolve. Once dissolved, this will yield a working solution containing **1000 ppm** of free chlorine dioxide. Prepare fresh solutions daily.

MB-10 TABLETS®

Disinfectant / Virucidal* / Tuberculocidal / Sterilant Food-Contact Surface Sanitizer / Oxidizing Tablets

For use in Hospitals, Medical, Dental, Industrial,
Manufacturing and Institutional Facilities, Laboratory
Animal Facilities, Clinical and Research Laboratories,
Veterinary Hospitals and Clinics, and Animal Rearing and
Confinement Facilities.

Active Ingredients:

Sodium Chlorite	20.8%
Sodium Dichloroisocyanurate dihydrate	7.0%
<u>Other Ingredients</u>	<u>72.2%</u>
Total	100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

See right side panel for First Aid and Precautionary Statements.

EPA Reg. No. 70060-19-46269 EPA Est. No. 46269-DE-01

NET WEIGHT:

1.5 Gram Tablets (150 ea.) 0.55 lbs. (225 grams)

Expiration Date: December 2008

Manufactured for:



Quip Laboratories, Inc.

1500 Eastlawn Avenue • Wilmington, DE 19802

TEL 800-424-2436 • FAX (302) 761-2611

Patent: 6,699,404

Contains



Aseptrol and the Aseptrol logo are trademarks of BASF Corporation.

Sodium Chlorite in this product is from Spain.

LABORATORY ANIMAL FACILITIES, ANIMAL REARING FACILITIES AND ANIMAL RESEARCH FACILITIES AND LABORATORIES

Use **MB-10 Tablets**® to disinfect and kill tuberculosis bacteria, HIV-1, Human coronavirus, Hepatitis B Virus, Hepatitis C Virus, Newcastle Disease virus, Norovirus, Foot and Mouth Disease Virus, Human Influenza A virus (Hong Kong), Avian Influenza A (H3N2) virus and animal viruses* on hard non-porous surfaces such as floors, walls, counters, stainless steel environmental surfaces, bio-safety hoods, sinks, tiles, cages, coops, crates, kennels, instruments and utensils. Pre-clean surfaces and then apply either a 100 ppm chlorine dioxide solution (for Human Influenza A virus (Hong Kong), Avian Influenza A (H3N2) virus and animal viruses*) or 200 ppm chlorine dioxide solution (for disinfection, HIV-1, Human coronavirus and TB control). Apply the use-solution with a cloth, mop, sponge or sprayer or by immersion. Treated surfaces must remain wet for 10 minutes (30 minutes for Foot and Mouth Disease Virus). Wipe dry with a cloth, sponge or mop or allow to air dry. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface and spray until totally wet. Do not breathe spray. Allow to air dry.

ANIMAL ROOM DISINFECTION DIRECTIONS USING AN ULTRA LOW VOLUME FOGGING DEVICE

Remove all animals and feed from animal room, vehicles and enclosures. Remove all litter from floors, walls and surfaces of the room to be treated. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Close room off so fog is confined to room to be treated. Mix two (2) 6 gram **MB-10 Tablets**® or eight (8) 1.5 gram **MB-10 Tablets**® into one gallon of water making a 200 ppm solution of chlorine dioxide. Place Ultra Low Volume (ULV) fogger in center of room or insert the nozzle of the fogger through a suitable opening into the room. With the Flow Rate setting in HIGH output, apply fog for 15 minutes for each 3000 cubic feet of space in the room, thorough wetting of all surfaces is required.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Do not allow people to enter treated room until ten air exchanges or 2 hours of mechanical ventilation (i.e., fans). If the building must be entered, then the individuals entering the building must wear a self contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING FOR ROOM AND MACHINE SURFACES.

HEALTH-CARE and VETERINARY FACILITIES

Use **MB-10 Tablets**® to disinfectant pre-cleaned surfaces and to decontaminate instruments in hospitals, medical and dental offices, veterinary offices and clinics and related facilities. Apply a 200 ppm chlorine dioxide use-solution to hard, non-porous surfaces and/or instruments thoroughly wetting surfaces with a cloth, mop, sponge or sprayer or by immersion. Treated surfaces must remain wet for 10 minutes (30 minutes for Foot and Mouth Disease Virus). Wipe dry with a cloth, sponge or mop or allow to air dry. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface and spray until totally wet. Do not breathe spray. Allow to air dry. The 200 ppm chlorine dioxide use-solution is effective against gram negative and gram positive bacteria, HIV-1, Human coronavirus, Hepatitis B Virus, Hepatitis C Virus, Newcastle Disease virus, Norovirus, Foot and Mouth Disease Virus, Human Influenza A virus (Hong Kong), Avian Influenza A (H3N2) virus, animal viruses*, and tuberculosis bacteria.

This product is not to be used as a terminal sterilant / high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to decontaminate pre-cleaned critical or semi-critical medical devices prior to sterilization or high-level disinfection.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV, HBV & HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS that involve healthcare settings, or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects soiled with blood or body fluids can be associated with the transmission of HIV-1, HBV and HCV. **MB-10 Tablets**® destroys HIV-1, HBV and HCV on pre-cleaned environmental surfaces/objects previously soiled with blood or other body fluids at 200 ppm and 1 minute contact (10 minutes for HBV and HCV).

PERSONAL PROTECTION: The worker should wear disposable latex gloves, gown, mask and eye protection to prevent contamination from soiled items.

CLEANING PROCEDURE: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of **MB-10 Tablets**® solution.

CONTACT TIME Allow **MB-10 Tablets**® solution to contact treated items for 1 minute (10 minutes for HBV and HCV). A contact time of 10 minutes will not control other common types of viruses and bacteria.

DISPOSAL OF INFECTIOUS MATERIALS: Any blood and other body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

SURFACE STERILIZATION

Use **MB-10 Tablets**® where sterility conditions are critical for optimum performance, such as manufacturing and laboratory equipment, and in areas where sterilization is required, such as Level 3 and 4 Biosafety Level (BSL-3 and BSL-4) facilities. Use **MB-10 Tablets**® on hard, non-porous surfaces such as plastics (polystyrene, polypropylene, polyvinyl chlorides, polyesters), stainless steel, fiberglass, ceramic, metal or glass. Do not use **MB-10 Tablets**® as a terminal high-level disinfectant or sterilant on any critical/semi-critical medical device or instrument. Prior to use, thoroughly pre-clean surface to be sterilized. This can be accomplished by rinsing with purified water, mechanical action or by detergent cleaning followed by a water rinse. Pre-cleaned surfaces may be allowed to air dry or may be towel dried but do not dry surfaces using dry heat. Prepare a 1000 ppm use-solution of chlorine dioxide by following the instructions under "Preparation of Use Solution" on this label. Apply the 1000 ppm chlorine dioxide use-solution by either thoroughly soaking the target surface or by immersion. All target surfaces must be exposed to treatment solution for at least 1 hour. Allow to air dry.

FOOD-CONTACT SURFACE SANITIZER

Use **MB-10 Tablets**® to sanitize hard, nonporous food-contact surfaces and utensils in food processing plants, breweries, bottling plants, restaurants and other food handling establishments. Prior to application, remove gross food particles and soil by a pre-flush, or pre-scrape and, when necessary, pre soak. Then thoroughly wash or flush surfaces with a good detergent or compatible cleaner followed by a potable water rinse before application of the sanitizer solution. Apply a use solution of 25 ppm (50 ppm for Listeria monocytogenes) chlorine dioxide to pre-cleaned hard surfaces thoroughly wetting surfaces with a cloth, mop, sponge, coarse sprayer or by immersion. Surfaces must remain wet for at least 60 seconds and then followed by adequate draining and air drying. Do not rinse. Prepare a fresh solution for each use.

SPECIAL INSTRUCTIONS FOR CLEANING AND DISINFECTING AREAS WHICH MAY BE INFESTED WITH HANTAVIRUS

Infection with Hantavirus occurs by inhalation of infectious materials. Persons involved in the clean-up must wear coveralls (disposable, if possible), rubber boots or disposable shoe covers, rubber or plastic gloves, protective goggles, and an appropriate respiratory protection device, such as a half-mask air-purifying (or negative-pressure) respirator with a high-efficiency particulate air (HEPA) filter or a powered air-purifying respirator (PAPR) with HEPA filters.

All potential infective waste material (including respirator filters) from clean-up operations that cannot be burned or deep buried on site must be double bagged in appropriate plastic bags. The bagged material must then be labeled as infectious (if it is to be transported) and disposed of in accordance with local requirements for infectious waste.

Rodent droppings and visible dust may be reservoirs for Hantavirus. If you are cleaning out a building that has been closed up, such as a cabin, shed, or garage:

- A. Air out building for at least 30 minutes by opening windows and doors.
- B. Leave the building while it is airing out.
- C. Do not vacuum, sweep or dust. This may spread the virus through the air.
- D. Thoroughly wet the contaminated areas with **MB-10 Tablets**® and allow to stand undisturbed for 10 minutes.
- E. Carefully remove non-salvageable contaminated material and dispose by burial or burning. Contact your local and state health department for additional disposal methods.
- F. Treat the surface again following the label directions and allow to stand undisturbed for 10 minutes.

For additional guidance visit CDC website at www.cdc.gov/hantavirus.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals**

DANGER. Corrosive. Causes irreversible eye damage and skin burns. May be fatal if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses), protective clothing and rubber gloves. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms.

STORAGE AND DISPOSAL

Store this product in a cool, dry area away from direct sunlight and heat to avoid deterioration and in an area inaccessible to children. Place empty pouch or blister pack and container in plastic bag and discard in trash.

Viruses:	Use-Dilution (ppm)	Time (minutes)
Minute Virus of Mouse (Parvovirus) (MVM-p)	100	10
Minute Virus of Mouse (Parvovirus) (MVM-H)	100	10
Mouse Hepatitis Virus (MHV-A59)	100	10
Mouse Hepatitis Virus (MHV-JHM)	100	10
Murine Parainfluenza Virus Type 1 (Sendai)	100	10
Sialodacryoadenitis Virus (Coronavirus) (SDAV)	100	10
Theiler's Mouse Encephalomyelitis Virus (TMEV)	100	10
Avian Influenza A (H3N2) virus	100	10
Canine Parvovirus	100	10
Foot & Mouth Disease virus	100 & 200	30
Hantavirus	100	10
Hepatitis B Virus	200	10
Hepatitis C Virus	200	10
HIV	200	1
Human Coronavirus	200	10
Infectious bursal disease virus (IBDV)	100	10
Influenza A virus (Hong Kong)	100	10
Marek's disease virus (MDV)	100	10
Newcastle Disease virus	200	5
Norovirus	200	10
Porcine circovirus type 2	100	10
Porcine Respiratory and Reproductive Syndrome virus (PRRSV)	100	10
Vaccinia virus	50 & 100	10
Bacteria:		
Bordetella bronchiseptica	200	10
Clostridium difficile (vegetative)	200	10
Corynebacterium bovis	100	10
Enterococcus faecalis Vancomycin Resistant	200	5
Escherichia coli O157:H7	200	10
Helicobacter pylori	100	10
Mycobacterium bovis	200	5
Pseudomonas aeruginosa	200	10
Salmonella enterica	200	10
Staphylococcus aureus	200	10
Staphylococcus aureus (MRSA)	100 & 200	10
Staphylococcus aureus Methicillin Resistant	200	5